International Journal of Advance Research in Community Health Nursing

E-ISSN: 2664-1666 P-ISSN: 2664-1658 IJARCHN 2019; 1(1): 16-18 Received: 19-11-2018 Accepted: 21-12-2018

Shubhangi Borude

Clinical Instructor, Bharati Vidyapeeth College of Nursing, Pune, Maharashtra, India

Sourabh More

Third Year GNM Nursing, Bharati Vidyapeeth School of Nursing, Pune, Maharashtra, India

Aishwarya Sanas

Third Year GNM Nursing, Bharati Vidyapeeth School of Nursing, Pune, Maharashtra, India

Rani Pandhare

Third Year GNM Nursing, Bharati Vidyapeeth School of Nursing, Pune, Maharashtra, India

Correspondence Shubhangi Borude Clinical Instructor, Bharati Vidyapeeth College of Nursing, Pune, Maharashtra, India

A study to assess the knowledge regarding selected water borne diseases among the people residing in selected areas of Pune

Shubhangi Borude, Sourabh More, Aishwarya Sanas and Rani Pandhare

Abstract

A study was undertaken to assess the knowledge regarding selected water borne diseases among the people residing in selected areas of Pune" to assess the knowledge regarding selected water borne diseases among the people To associates the findings with selected demographic variables.

The objectives of the study were to assess the knowledge regarding selected water borne diseases among the people To associates the findings with selected demographic variables Quantitative approach was chosen for the research study. The research design used for the study was exploratory survey design. The study consists of 200 samples which were selected by non probability convenient sampling. The study was conducted in Singhgad Road, Hingne Khurad, Pune Guruwar Peth, Swarget, pune. Katraj gaonthan, Pune. Hadapsar, pune, Pune. The data was collected by administering structured questionnaire to 200 participants under study. The questionnaire consists of two parts. Section-A which deals with demographic data of participant and Section- B which consist of questions related to knowledge regarding selected water borne diseases.

The validity of the tool was done by 5 experts from various fields of Nursing i.e, Community helath nursing, Obstetrics and Gynecology Nursing, Medical and Surgical Nursing. The reliability of questionnaire was established by the method of test retest method and was found to be 0.89.

In these study it is found that most people were having average knowledge but still there are people whose knowledge level are poor. It shows that water borne diseases are still limited and it requires greater attention commitment from health professional. There is significant association between the selected demographic variables and the knowledge of the selected water borne diseases i.e., age. Recommendations were made based on the findings of the study. Researcher recommends that a similar study can be replicated in different setting to strengthen the findings, the same study can be replicated on a larger sample for each group for the generalization of the findings, A comparative study can be conducted on knowledge regarding selected water borne diseases after a teaching programme on prevention and control of water borne diseases to people.

Keywords: Knowledge, water borne diseases, people

1. Introduction

Water is the most indispensable nature resource in the world for every living being. The entire life-support systems are dependent upon this vital resource. It is most important to all as it is directly consumed by all living entities. From that point of view, water is particularly related with health. Ninety percent of cases are reportedly attributed to the supply of unsafe drinking water coupled with improper sanitation and poor hygiene.

Adequate supply of fresh and clean drinking water is a basic need for all human beings on the earth, yet it has been observed that millions of people worldwide are deprived of this. Industrial growth, urbanization and the increasing use of synthetic organic substances have serious and adverse impacts on freshwater bodies. Many areas of groundwater and surface water are now contaminated with heavy metals, POPs (persistent organic pollutants), and nutrients that have an adverse affect on health The burden of waterborne diseases is paramount in the globe. About 4% of the global burden of diseases is attributable to water, sanitation and hygiene. Nearly 2.2 million people die every year due to diarrhoeal diseases globally. Of these, 1.8 million deaths occur alone in low-income countries. Further, in low and middle-income countries one of the tenth leading causes of death is attributable to diarrhoea-related diseases. Globally, diarrhoea alone kills more children compared to malaria and tuberculosis together,

The awareness of peoples about waterborne disease and preventive services is a barometer by which we can measure the progress of family, community and country. Lack of awareness can lead to health hazards in country.

2. Methodology

A non experimental research design was adopted to conduct the study. A total of 200 samples was selected by using non probability convenient sampling technique. Study instrument was self-structured questionnaire used by the researcher, section 1 consist of demographic variables and section 2 consist of structured knowledge questionnaire. The tool was content validated by expert and translated in to Marathi which was again validate. The value is 0.89 which is highly reliable The samples were selected by using non probability convenience sampling method. The collected data were analyzed by using descriptive and inferential statistics.

3. Results

Table 1: Demographic description of sample by frequency and percentage of demographic data N=200

| Sr. no | Characteristics | Frequency | Percentage |
|--------|------------------------|-----------|------------|
| | Age | | |
| | a) 14 to 18 years | 73 | 19.5% |
| 1. | b) 19 to 30 years | 66 | 36.5% |
| | c) 31 to 45 years | 39 | 33% |
| | d) 46 to 60 years | 22 | 11% |
| | Religion | | |
| | a) Hindu | 173 | 86.5% |
| 2 | b) Muslim | 09 | 4.5% |
| | c) Christian | 03 | 1.5% |
| | d) Any other | 15 | 7.5% |
| | Education | | |
| | a) Primary education | 27 | 13.5% |
| 2 | b) Secondary education | 109 | 54.5% |
| 3 | c) Graduate | 51 | 25.5% |
| | d) Post graduate | 11 | 5.5% |
| | e) Illiterate | 02 | 01% |
| | Occupation | | |
| | a) Housemaker | 38 | 19% |
| | b) Job | 75 | 37.5% |
| 4 | c) Business | 27 | 13.5% |
| | d) Worker | 33 | 16.5% |
| | e) Student | 27 | 13.5% |
| | Family income | | |
| | a) Below Rs 10000 | 38 | 19% |
| 5 | b) Rs 10001 to 20000 | 75 | 37.5% |
| | c) Rs 20001 to 40000 | 54 | 27% |
| | d) Above Rs 40001 | 33 | 16.5% |
| | Family type | | |
| | a) Joint family | 81 | 40.5% |
| 6. | b) Nuclear family | 103 | 51.5% |
| | c) Extended family | 10 | 05% |
| | d) Any other | 06 | 03% |

The data given in table-1 shows that Maximum sample (36.5%) were in the age group of 19 to 30 years. Maximum of samples (86.5%) were Hindu. Maximum sample (54.5%) studied till Secondary education. Maximum sample (37.5%) are having job as a Occupation. Maximum samples (37.5%) are having family income between Rs 10001-20000.Maximum samples (51.5%) belongs to nuclear family Maximum (49.5%)

Table 2: Showing knowledge score of knowledge regarding selected water borne diseases among the people of selected area of Pune. N=200

| Sr. No | Level of knowledge | frequency | Percentage |
|--------|-----------------------------|-----------|------------|
| 1 | 00-09 (poor knowledge) | 20 | 10% |
| 2 | 10-20 (average knowledge) | 99 | 49.5% |
| 3 | 21-25 (good knowledge) | 71 | 35.5% |
| 4 | 26-30 (Excellent knowledge) | 10 | 5% |

The above table shows that out of samples,49.5% of the people are showing the average knowledge about selected water born disease, 10% are showing poor knowledge about selected water borne disease.

Table 3: Mean and standard deviation of the knowledge of postnatal mothers regarding self-care after childbirth, N=100

| Sr. No | Categories | Mean N=200 | Standard deviation N=200 |
|-----------|----------------------|---------------|-----------------------------|
| 1 | Mean knowledge score | 18.135 | 0.97858 |

The data represents in table-3 shows that the mean and standard deviation is 18.135 and 0.2390 respectively.

Table 4: Association of knowledge score with demographic data

| Demographic | Degree of | Table | Calculated | P- value | |
|---------------|-------------|-------|------------|-----------|--|
| data | association | value | value | r - value | |
| Age | 9 | 23.36 | 16.92 | 0.01 | |
| Religion | 9 | 13.06 | 14.68 | 0.1 | |
| Education | 12 | 15.24 | 18.55 | 0.90 | |
| Occupation | 12 | 7.368 | 18.55 | 0.95 | |
| Family income | 9 | 10.62 | 14.68 | 0.1 | |
| Family type | 9 | 8.890 | 14.68 | 0.90 | |

The above table depletes that there is a association of Age with knowledge as the calculated chi square value is (23.26) greater than table value (16.92), And as the p value is less than 0.05

5. Conclusion

The purpose of the present study was to assess the knowledge regarding selected water borne diseases among the selected area of pune. The 200 sample were selected from selected area of pune city i.e, Singhgadroad, Hingane Khrud, Pune and other selected areas. Descriptive research design was used for the study. The content validity and reliability of the tool was done, which suggested that the tool was reliable. The pilot study was conducted on 20 samples and the feasibility of the study was established. It was found that the tool had no major flaws and was used for the final study with the changes as per the experts and Guide.

Based on the objectives, the collected data was analyzed by using descriptive statistics.

In this majority of the samples i.e49.50% of samples having average knowledge regarding selected water borne diseases 35.50% having good knowledge and 4% sample shows excellent knowledge and10% sample having poor knowledge category.

6. Recommendation

- A similar study can be replicated in different setting to strengthen the findings.
- The same study could be replicated on a large sample.
 This would provide invaluable evidence in the area of practice.
- A comparative study can be conducted of the pre and

- post test knowledge of the selected water borne diseases after a teaching programme on prevention and control of water borne diseases to people.
- A comparative study can be conducted on knowledge regarding selected water borne diseases after a teaching programme on prevention and control of water borne diseases to people.
- The study can be done on association between various demographic variables, which were significant, on larger samples.
- A follow up study can be conducted to evaluate effectiveness of planned teaching programme on water borne diseases among people.

7. References

- 1. "http://en.wikipedia.org/wiki/ Water-
- 2. Narongsak Noosorn, prasit. Journal of American Water Works Association. 2000; 9:24-43.
- 3. Kamala *et al.* Journal of water for people, water for life, The united nations world water. Development Report. 2001; 4:102-105
- 4. Khamgaonkar *et al.* Journal of medical science, Awareness of mothers regarding waterborne diseases, 6(44):55-61.
- 5. Mittal SK. Indian Journal of Paediatrics, Diarrhoea In Tropics. 1999; 65(4):4-5.
- 6. Aidea. global (n.d) Waterborne Diseases (Electronic Version). Retrieved 21st 2009, from http://www.aldeaglobal.com/ar/agua/wbd.htm.
- 7. Shruff. Journal of Public Health and Epidemiology, Public Participation to minimize Waterborne diseases in Yom riverside communities. 2005; 1(1):09-23.
- 8. Dua *et al.* Indian Journal of paediatrics, prevalence of Diarrhoea, 9:55-59.